



Bluetongue and epizootic hemorrhagic disease viruses: recent developments with these globally re-emerging arboviral infections of ruminants

Nigel James Maclachlan¹, Stephan Zientara², William C Wilson³, Juergen A Richt⁴ and Giovanni Savini⁵

Bluetongue (BT) and epizootic hemorrhagic disease (EHD) are globally re-emerging diseases of domestic and wild ruminants, respectively caused by BT virus (BTV) and EHD virus. Both viruses are transmitted by hematophagous midges; however, newly recognized BTV serotypes may be transmitted horizontally without requirement for any biological vector. The global range of these viruses and/or their associated diseases have changed remarkably in recent years, most notably with the invasion of Europe by multiple serotypes of BTV since 1998. Although not zoonoses, the unanticipated emergence of BT and EHD in several different areas of the world provides a uniquely sobering and unambiguous reminder of the potential consequences of climate change on the distribution and severity of vector-borne diseases. Recent experiences with these viruses have also emphasized the need for effective, DIVA-compatible vaccines to combat anticipated future incursions, as existing vaccines have serious inherent deficiencies.

Addresses

¹ Department of Pathology, Microbiology and Immunology, School of Veterinary Medicine, University of California, Davis, CA 95616, USA

² UMR VIROLOGIE, INRA, Ecole Nationale Vétérinaire d'Alfort, ANSES, Université Paris-Est, Maisons-Alfort 94700, France

³ Arthropod-Borne Animal Diseases Research Unit, Agricultural Research Service, USDA, Manhattan, KS, USA

⁴ Department of Diagnostic Medicine/Pathobiology, College of Veterinary Medicine, Kansas State University, Manhattan, KS, USA

⁵ Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise, OIE Reference Laboratory for BTV, Teramo, Italy

Corresponding author:

Maclachlan, Nigel James (njmaclachlan@ucdavis.edu)

Current Opinion in Virology 2019, 34:56–62

This review comes from a themed issue on **Emerging viruses: inter-species transmission**

Edited by **Adolfo García-Sastre** and **Juergen A Richt**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 14th January 2019

<https://doi.org/10.1016/j.coviro.2018.12.005>

1879-6257/© 2018 Elsevier B.V. All rights reserved.

Introduction

Bluetongue virus (BTV) and epizootic hemorrhagic disease virus (EHDV) are closely related arboviruses that are classified in the genus *Orbivirus* within the subfamily Sedoreovirinae of the family Reoviridae [1*,2*,3]. Both viruses are transmitted to wild and domestic ruminants by certain species of hematophagous *Culicoides* midges. Whereas, BTV is the cause of significant economic loss to ruminant livestock production, especially of sheep, EHDV is most significant as a pathogen of ruminant wildlife, notably white-tailed deer (*Odocoileus virginianus*). BTV and EHDV are genetically diverse, with multiple serotypes and myriads of genetically distinct strains of each virus [4,5*,6,7*,8,9]. Genetic diversity within these virus species occurs as a consequence of both genetic drift and genetic shift; genetic shift occurs as a consequence of reassortment of viral gene segments during co-infections of either animals (ruminants) or insect vectors (*Culicoides* midges) with more than one virus serotype or strain, whereas genetic drift occurs as a consequence of polymerase ‘infidelity’ during replication of the virus leading to a viral quasispecies with subsequent founder effect of specific genetic variants in either insect or animal hosts. The individual species of *Culicoides* midge that serve as vectors of these viruses differ between regions of the world, as do the serotypes and strains of BTV and EHDV that occur within each global ‘episystem’ [4,9,10*,11]. Thus, the respective global ranges of BTV and EHDV infections are similar but certainly not identical. Furthermore, there have been profound recent changes in the global distribution and biological behavior of both BTV and EHDV. Given that the epidemiology, pathogenesis and other features of BTV and EHDV infections have recently been reviewed [1*,2*,3], the purpose of this article is to summarize recent novel developments regarding characterization of the global ecology and biological behavior of these two viruses.

Bluetongue

Expansion of global range

The global range of BTV infection has changed remarkably with the advent of the new millennium, notably at the traditional upper and lower limits of the virus’ global range (approximately 50° North and 35° South) [2*]. Climate change is proposed to have contributed this expansion in the traditional global range of BTV as a consequence of its impact on the distribution of vector-competent species of

midges as well as the vectorial capacity of populations of midges already resident in different regions of the world (including those not previously thought to be competent vectors) [12[•]]. Notable recent changes in the global distribution of BTV include: 1. Incursion of multiple serotypes of BTV into Europe [2[•],12[•]]. Before 1998, only portions of Europe within the Mediterranean Basin were subject to incursions of BTV, and these were transient and involved only single virus serotypes; 2. Incursion since 1999 of multiple novel BTV serotypes into the southeastern United States from the adjacent Caribbean episystem [2[•],13]; 3. First detection of BTV in Ontario, Canada, in 2015 as a result of northern spread of the virus from the United States [14]; 4. Extension of BTV into Victoria, Australia, in 2017, which is farther south than the virus' traditional distribution on that continent [15]; 5. Increasing detection of multiple BTV serotypes in South America and Asia, including countries such as South Korea that had not previously documented such infections [16–18].


Notable recent findings regarding the global epidemiology of BTV infection include (see Figure 1):

- 1 *Reemergence of BTV serotype 8 in France.* BTV serotype 8 (BTV-8) first emerged in northern Europe in 2006, although the origin and route of introduction of the particular strain of BTV-8 remain uncertain [2[•]]. BTV-8 spread rapidly throughout northern Europe in species of *Culicoides* midge not previously known to be efficient vectors of BTV. After an intensive vaccination campaign against BTV-8, France and adjacent countries of northern Europe were declared free of BTV as of December 2012. France also was declared free of BTV-1, which spread from the Mediterranean region to northern France in 2007. However, surveillance has confirmed that BTV-8 has continued to circulate 'silently' among livestock in France [19,20,21[•]]. Interestingly, although the original (2006) and recent (2015) strains of BTV-8 are genetically similar or identical, recent infections have invariably been subclinical, whereas clinical disease was common among ruminant livestock (including cattle) infected during the original outbreak.
- 2 *BTV-4 in France.* BTV-4 has been detected frequently in the Mediterranean region since 1998, and again appeared in Corsica in 2016, likely as a result of spread from adjacent infected Mediterranean countries where the virus continues to circulate [22]. A genetically distinct reassortant strain of BTV-4 that was identified in the Balkan countries (2014), mainland Italy (2014–2016), and Sardinia (2016–17) was identified recently in a healthy calf in mainland France (adjacent to the Swiss border) [23].
- 3 *BTV-3 and BTV-6 in the Mediterranean Basin.* More than 60 years after its original and only detection in Cyprus, BTV-3 reappeared in the Mediterranean Basin causing disease among sheep in Israel in 2013 and 2016–17. Similar strains of BTV-3 were also identified in Tunisia and Italy, consistent with spread of the virus via infective vectors or illegal animal movement from North Africa to Italy [24[•]]. These strains of BTV-3 were associated with typical signs of bluetongue among sheep in both regions, but not in cattle. In contrast, BTV-6 also reemerged in Israel in 2017 but, unlike BTV-3, BTV-6 infection resulted in clinical disease in both cattle and sheep throughout most of Israel (G Savini, unpublished data).
- 4 *BTV-3 in the United States.* BTV-10, 11, 13, and 17 have long been endemic in much of the United States, whereas BTV-2 is limited to the Southeastern United States and, recently, California [2[•],7[•],13]. Although some 11 previously exotic serotypes of BTV have invaded the Southeastern United States from the Caribbean Basin episystem since 1999, most of these viruses have not spread far from their original site of incursion. In contrast, BTV-3 has spread widely since 2006, and it is now clearly evident that the emergent strains of BTV-3 have extensively reassorted genes with those of the traditional endemic serotypes (BTV-10, 11, 13,17).
- 5 *Live-attenuated BTV-14 vaccine in Europe.* At least 3 (BTV-6, BTV-11, BTV-14) different South African live-attenuated BTV vaccines have been disseminated by vectors among livestock in Europe, most recently BTV-14 in western Russia, Poland, Lithuania and Spain (in a single animal imported from Lithuania) [25,26,27[•]]. It remains uncertain how these viruses were introduced into Europe as they were not used in official vaccination campaigns. The natural spread of live-attenuated BTV vaccine viruses also occurs elsewhere in the world, and genetic analyses confirm that these vaccine BTVs can reassort their genes with those of other endemic BTVs to create genetically distinct progeny, potentially with novel biological properties [2[•],28[•]].

Identification of novel, small ruminant adapted BTVs

Several new serotypes of BTV have been described recently in Europe, Asia and Africa. These viruses represent a novel group of BTVs that differ in their biological and genetic properties from the historical 24 serotypes (BTV 1–24) that previously have been identified globally. Specifically, these novel BTVs exhibit minimal virulence or pathogenicity to ruminants and have all been isolated from apparently healthy goats or sheep. Some of these viruses can be transmitted by contact (horizontally) without requirement for vector insects, and some appear to cause persistent infections in their animal hosts. Although little is known about these new BTV types, some have already been classified as additional serotypes (specifically BTV 25–27), whereas others have yet to be assigned a numerical serotype designation. Serological classification of some of

Figure 1



Recently described changes in the global distribution or biology of bluetongue virus (BTV) depicting affected countries (not the precise distribution of each virus). **(a)** 'Traditional' serotypes of BTV (1–24) in Europe and the Mediterranean Basin (additional serotypes circulating are noted directly on the affected areas). Adapted from a map showing restriction zones for bluetongue virus in the European Union as of May 30, 2018 (available at http://www.bluetonguevirus.org/sites/blue/files/quick_media/Current%20EU%20BTV%20Restriction%20Zones.jpg). **(b)** Spread of BTV-3 from the southeastern United States. **(c)** Identification of new serotypes (BTV 25–27) and related novel viruses (BTV X, Y and SP) in Europe, Africa and the Middle East (BTV XJ1407 from China is not shown).

these novel viruses is complicated because of their resistance to propagation in cell culture.

The first of these novel BTVs was named Toggenburg Orbivirus after the region of its first detection in goats in Switzerland. Sequence analyses confirmed the virus to be a new BTV serotype, namely BTV-25 [29,30]. Although first detected in 2008, retrospective analyses confirmed

positive samples from Swiss goats dating back to 1998 indicating that BTV-25 has circulated in that part of Europe for at least a decade and likely considerably longer. BTV-25 produces subclinical infections in goats and is only mildly pathogenic to sheep. Although BTV-25 infection is highly prevalent among goats in certain areas of Switzerland, infection was not found in either sheep or cattle cohabiting with virus-positive goats, nor in wild

animals, suggesting a highly restricted host range [31]. The high prevalence of BTV-25 infection in goats likely reflects its ability to be transmitted horizontally and induce persistent infection in this species. Experimental studies are difficult with BTV-25 because the virus is yet to be propagated *in vitro* using embryonated chicken eggs (ECE) or cell culture systems that are reliably used for propagation of BTV 1–24 [32]. A probable variant of BTV-25, BTV-Z ITA2017, was recently identified in a goat flock from North Western Italy [33]. Like BTV-25, ruminants infected with BTV-Z ITA2017 showed no clinical signs and only low levels of RNA-emia.

Another novel BTV, BTV-26, was first isolated in 2010 from the blood of a sheep in Kuwait [34]. Blood samples were positive by BTV group-specific RT-PCR, but consistently negative in serotype-specific RT-PCR assays, indicating the presence of a novel BTV serotype. The sequence of the serotype-specific VP2 gene of BTV-26 differs from other known serotypes (BTV 1–25), but is more closely related to BTV-25 than other serotypes. Unlike BTV-25, BTV-26 can be propagated *in vitro* although its growth capability is limited to mammalian cell lines and the virus does not replicate in culicoides-derived cells (Kc). Experimental BTV-26 infection results in a higher and longer-lasting RNA-emia in goats than in sheep. Whereas BTV-26 infection is subclinical in goats, the virus causes mild disease in sheep [35,36]. BTV-26 can be spread between goats by direct contact transmission via either nasal or ocular secretions. Serological evidence of BTV-26 infection subsequently was shown among cattle and dromedaries in Mauritania and in Libyan sheep [37].

A further novel BTV serotype, now designated BTV-27, was first detected among goats in Corsica in 2014 [38]. The infected goats were positive by BTV group-specific PCR but negative in serotype-specific RT-PCR assays to each of the other 26 BTV serotypes. Sequence analyses confirm high identity with both BTV-25 and BTV-26; however, BTV-27 is not neutralized by antisera raised against other BTV serotypes. Like BTV-26, *in vitro* propagation of BTV-27 is limited to selected cell lines; specifically, the virus does not grow in ECE or in cell cultures (e.g. Kc, Vero and BHK) that are commonly used for isolation of 'traditional' BTV serotypes. Several genetically related but distinct strains of BTV-27 have been identified in healthy goats, whereas cattle appear to be refractory to infection. Like other small ruminant adapted BTVs, BTV-27 can be spread horizontally by direct contact transmission [39].

A number of other genetically distinct BTVs recently have been identified that likely represent additional virus serotypes. These include a virus designated X ITL2015 from healthy goats in Italy [40]. Although clearly a BTV in terms of reaction with group-specific RT-PCR and BTV-group specific serological testing of infected animals, the virus is virologically and serologically distinct from BTV

1–27 and is genetically most similar to recently isolated strains of BTV-27 from Corsica and to another recently discovered BTV strain (XJ1407) from China. Efforts to date to isolate X ITL2015 using ECE or cell cultures have been unsuccessful. The virus designated XJ1407 was first detected in clinically normal goats and sheep in China [41]. Although it is most related to BTV X ITL2015, and less so to BTV-25 and 27, XJ1407 can be propagated in ECE and several mammalian and insect cell lines.

Sheeppox vaccine-derived BTV (SPV BTV) was identified in 2014 in batches of a sheeppox vaccine used in Israel [42]. This virus is closely related to BTV-26 and, like BTV-26, the virus is easily propagated in ECE and several mammalian cell lines. A genetically similar virus (BTV-Y TUN2017) was identified among healthy sheep in North Africa (Tunisia) [24]. Unlike SPV BTV, however, BTV-Y TUN2017 does not grow in ECE or cell culture. SPV and related BTVs are only distantly related to BTV 1–27, and serum from BTV-Y TUN2017 infected sheep failed to neutralize other BTV serotypes, suggesting that it is a novel serotype. As the small ruminant adapted BTVs cause subclinical or very mild infections and their presence currently does not lead to restrictions on animal movement or trade (in contrast to those associated with the presence of BTV 1–24), these viruses currently are not problematic to the livestock industries of countries in which they are endemic. However, the biological characteristics of the newly recognized BTV serotypes are remarkably different from those of the 'classical' serotypes, and reassortment events between them could potentially result in progeny viruses with different properties [40,42,43].

BTV infections of dogs

The initial description of BTV infection of dogs resulted from problems associated with the widespread use in the United States of a live-attenuated canine vaccine contaminated with BTV-11 [44,45]. More recent canine BTV-11 infections in the United States were not associated with the use of contaminated vaccines, and the canine viruses were genetically similar to those circulating in livestock at the same time in the same area [46,47]. Similarly, BTV infection of dogs is prevalent in North Africa, and is associated with neither contaminated canine vaccines nor the ingestion of BTV-infected meat [48]. It is assumed, therefore, that dogs may become infected through the bites of BTV-infected vector midges, as also has recently been proposed for related African horse sickness virus infections of dogs in South Africa [49]. BTV-11 infection is especially devastating in pregnant bitches, leading to abortion and death of the affected bitch.

The phenomenon of BTV over-wintering

Infection with traditional BTV serotypes (BTV 1–24) is seasonal in temperate regions of the world, reflecting the abundance of vector midges during late summer and autumn. How BTV persists between seasons in such

regions has been a topic of considerable controversy as transovarial transmission of virus has not been described in *Culicoides* spp. [50]. Recent studies in BTV-endemic temperate regions confirm the presence of low numbers of BTV-infected adult midges during winter months, suggesting that BTV can over-winter in long-lived vectors [51–53]. Of particular relevance is the fact that BTV-infected midges display profoundly altered feeding behavior that results in markedly reduced attraction to the light traps that have long been the mainstay of vector surveillance for BTV. As a result, exclusive use of light traps for vector surveillance can result in serious errors in determination of the regional risk of BTV transmission, and failure to detect the presence of BTV-infected, over-wintering midges [54*].

Epizootic hemorrhagic disease

Epizootic hemorrhagic disease (EHD) is a significant cause of both morbidity and mortality in North American white-tailed deer and certain other species of wild ruminants whereas, with the notable exception of Ibaraki virus, EHDV infections of ruminant livestock are typically subclinical [1*,3]. There are 7 serotypes of EHDV recognized currently, and Ibaraki virus, which is the cause of sporadic disease outbreaks in cattle in Asia, is a representative of serotype 2 (EHDV-2). Although the global distribution and epidemiology of BTV and EHDV infections are similar, EHDV infections have been less intensively studied because they are of less significance to animal agriculture as compared to those caused by BTV, particularly in regards to restrictive non-tariff trade barriers imposed on livestock from BTV endemic areas. Although most natural and experimental EHDV infections (other than with Ibaraki virus) of livestock are subclinical or asymptomatic, outbreaks of EHD recently have been reported among cattle in the Mediterranean Basin, Reunion Island, South Africa, the United States, and elsewhere [1*,3,16,55–57]. It is clearly apparent that certain strains of EHDV are more virulent to cattle than others, notably Ibaraki virus (EHDV-2) and the recently described strains of EHDV-2, 6, and 7 that caused disease among cattle in several countries. Like BTV, however, the molecular basis of EHDV virulence is uncharacterized as other strains of these same EHDV serotypes (EHDV-2,6,7) typically do not cause significant disease among cattle in other endemic regions of the world. Similarly, whole genome sequence comparisons showed no obvious differences in the strains of EHDV 1 and 2 associated with disease outbreaks among cattle in the Midwestern United States and viruses from apparently unaffected animals in the same region [7*,8] (WC Wilson, unpublished data). The role of host and the related environmental factors in precipitating outbreaks of EHD in cattle are poorly defined.

There is considerable genetic variation among the strains of EHDV that circulate within endemic regions, including virus strains of the same serotype. For example, at

least three different serotypes of EHDV currently circulate in North America, one of which (EHDV-6) is a relatively recently introduced serotype. EHDV-6 spread into the Southeastern United States from the adjacent Caribbean ecosystem but, like BTV-3, only strains of EHDV-6 that include reassorted genes from the historically endemic EHDVs (EHDV-1,2) were able to expand their range beyond the southeastern United States.

Lessons learned from the global emergence/re-emergence of BTV and EHDV

- 1 It has long been speculated that climate change will lead to expansion of the global range of specific arboviruses and their associated diseases, but recent changes in the global distribution and impacts of both BTV and EHDV infections of ruminants have been especially dramatic. Not only has climate change apparently lead to changes in the dynamics of specific vectors, but it also has been associated with the transition of indigenous midges not thought to be virus vectors into ones that efficiently serve as biological vectors of BTV in particular.
- 2 The enormous genetic diversity of these orbiviruses (including anthropogenic influences such as use of live-attenuated vaccines) can lead to the emergence of viruses with unique biological properties, including the capacity for horizontal and vertical (transplacental) transmission and spread to other species (e.g. dogs). Reassortment of the genes of incursional orbiviruses with those already present (endemic) viruses has facilitated the successful introduction and spread of novel reassortant progeny between episytems.
- 3 Live-attenuated orbivirus vaccines clearly have serious inherent limitations, including their capacity for spread by vector midges, reassortment with endemic viruses to create genetically novel progeny, and expression of novel properties including transplacental transmission, transmission to heterologous species, etc. [2*,28*,58,59].
- 4 Novel vaccination strategies employing safe and effective DIVA-compatible vaccines will be needed to limit the future spread and economic impact of these re-emerging orbivirus infections of ruminants [2*,28*,60–62].

Conflicts of interest statement

Nothing declared.

Acknowledgements

Figures were generously provided by Dr Erin E. Schirtzinger, Arthropod-Borne Animal Diseases Research Unit, Agricultural Research Service, USDA, Manhattan, KS, USA.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest

1. Maclachlan NJ, Zientara S, Savini G, Daniels PW: **Epizootic haemorrhagic disease**. *Rev Sci Tech* 2015, **34**:341–351.

A comprehensive review of epizootic hemorrhagic disease and epizootic hemorrhagic disease virus, as of 2015.

2. Maclachlan NJ, Mayo CE, Daniels PW, Savini G, Zientara S, Gibbs EP: **Bluetongue**. *Rev Sci Tech* 2015, **34**:329-340.

A comprehensive review of bluetongue and bluetongue virus, as of 2015.

3. Savini G, Afonso A, Mellor P, Aradaib I, Yadin H, Sanaa M, Wilson W, Monaco F, Domingo M: **Epizootic haemorrhagic disease**. *Res Vet Sci* 2011, **91**:1-17.
4. Bonneau KR, MacLachlan NJ: **Genetic diversification of field strains of bluetongue virus**. *Vet Ital* 2004, **40**:446-447.
5. Firth C, Blasdel KR, Amos-Ritchie R, Sendow I, Agnihotri K, Boyle DB, Daniels P, Kirkland PD, Walker PJ: **Genomic analysis of bluetongue virus episystems in Australia and Indonesia**. *Vet Res* 2017, **48**:8.

This recent study details genetic characterization of bluetongue viruses from Australia and Indonesia, confirming the concept of regional co-evolution of virus strains and vectors in distinct global epidemiological systems (episystems).

6. Nomikou K, Hughes J, Wash R, Kellam P, Breard E, Zientara S, Palmarini M, Biek R, Mertens P: **Widespread reassortment shapes the evolution and epidemiology of bluetongue virus following European invasion**. *PLoS Pathog* 2015, **11**:e1005056.
7. Wilson WC, Gaudreault NN, Jasperson DC, Johnson DJ, Ostlund EN, Chase CL, Ruder MG, Stallknecht DE: **Molecular evolution of American field strains of bluetongue and epizootic hemorrhagic disease viruses**. *Vet Ital* 2015, **51**:269-273.

This study reports the genetic characterization of field strains of epizootic hemorrhagic disease and bluetongue viruses from different regions of the United States. A central conclusion is that the spread of novel serotypes of each virus that incurred originally into the southeastern United States was dependent on reassortment of genes of these incursional orbiviruses with those of viruses of other serotypes that were already present (endemic).

8. Wilson WC, Ruder MG, Jasperson D, Smith TP, Naraghi-Arani P, Lenhoff R, Stallknecht DE, Valdivia-Granda WA, Sheoran D: **Molecular evolution of epizootic hemorrhagic disease viruses in North America based on historical isolates using motif fingerprints**. *Virus Genes* 2016, **52**:495-508.
9. Balasuriya UB, Nadler SA, Wilson WC, Pritchard LI, Smythe AB, Savini G, Monaco F, De Santis P, Zhang N, Tabachnick WJ *et al.*: **The NS3 proteins of global strains of bluetongue virus evolve into regional topotypes through negative (purifying) selection**. *Vet Microbiol* 2008, **126**:91-100.
10. Gibbs EP, Greiner EC: **The epidemiology of bluetongue**. *Comp Immunol Microbiol Infect Dis* 1994, **17**:207-220.

This manuscript provides the initial proposal for global regionalization of bluetongue virus infection, specifically the concept of distinct global epidemiological systems (episystems).

11. Tabachnick WJ: **Culicoides and the global epidemiology of bluetongue virus infection**. *Vet Ital* 2004, **40**:144-150.
12. Baylis M, Caminade C, Turner J, Jones AE: **The role of climate change in a developing threat: the case of bluetongue in Europe**. *Rev Sci Tech* 2017, **36**:467-478.

A recent review of climatic factors as potential drivers of the emergence of bluetongue in Europe since 1998.

13. Schirtzinger EE, Jasperson DC, Ostlund EN, Johnson DJ, Wilson WC: **Recent US bluetongue virus serotype 3 isolates found outside of Florida indicate evidence of reassortment with co-circulating endemic serotypes**. *J Gen Virol* 2018, **99**:157-168.
14. Anon. *Canadian Food Inspection Agency*, available at <http://www.inspection.gc.ca/animals/terrestrial-animals/diseases/reportable/bluetongue/fact-sheet/eng/1306116803992/1306121522520>.
15. Anon. *Agriculture Victoria*, available at <http://agriculture.vic.gov.au/agriculture/pests-diseases-and-weeds/animal-diseases/general-livestock-diseases/bluetongue-virus>.
16. Verdezoto J, Breard E, Viarouge C, Quenault H, Lucas P, Sailleau C, Zientara S, Augot D, Zapata S: **Novel serotype of bluetongue virus in South America and first report of epizootic haemorrhagic disease virus in Ecuador**. *Transbound Emerg Dis* 2018, **65**:244-247.

17. Legisa DM, Gonzalez FN, Dus Santos MJ: **Bluetongue virus in South America, Central America and the Caribbean**. *Virus Res* 2014, **182**:87-94.
18. Seo HJ, Park JY, Cho YS, Cho IS, Yeh JY: **First report of Bluetongue virus isolation in the Republic of Korea and analysis of the complete coding sequence of the segment 2 gene**. *Virus Genes* 2015, **50**:156-159.
19. Bréard E, Sailleau C, Quenault H, Lucas P, Viarouge C, Touzain F, Fablet A, Vitour D, Attoui H, Zientara S *et al.*: **Complete genome sequence of Bluetongue virus serotype 8, which reemerged in France in August 2015**. *Genome Announc* 2016, **4**:e00163-16.
20. Courtejoie N, Durand B, Bournez L, Gorlier A, Bréard E, Sailleau C, Vitour D, Zientara S, Baurier F, Gourmelen C *et al.*: **Circulation of bluetongue virus 8 in French cattle, before and after the re-emergence in 2015**. *Transbound Emerg Dis* 2018, **65**:281-284.
21. Sailleau C, Bréard E, Viarouge C, Vitour D, Romey A, Garnier A, Fablet A, Lowenski S, Gorna K, Caignard G *et al.*: **Re-emergence of Bluetongue virus serotype 8 in France, 2015**. *Transbound Emerg Dis* 2017, **64**:998-1000.
22. Sailleau C, Breard E, Viarouge C, Gorlier A, Quenault H, Hirchaud E, Touzain F, Blanchard Y, Vitour D, Zientara S: **Complete genome sequence of bluetongue virus serotype 4 that emerged on the French island of Corsica in December 2016**. *Transbound Emerg Dis* 2018, **65**:e194-e197.
23. Sailleau C, Breard E, Viarouge C, Gorlier A, Leroux A, Hirchaud E, Lucas P, Blanchard Y, Vitour D, Grandcollet-Chabot M *et al.*: **Emergence of bluetongue virus serotype 4 in mainland France in November 2017**. *Transbound Emerg Dis* 2018, **65**:1158-1162.
24. Lorusso A, Sghaier S, Di Domenico M, Barbria ME, Zaccaria G, Megdich A, Portanti O, Seliman IB, Spedicato M, Pizzurro F *et al.*: **Analysis of bluetongue serotype 3 spread in Tunisia and discovery of a novel strain related to the bluetongue virus isolated from a commercial sheep pox vaccine**. *Infect Genet Evol* 2018, **59**:63-71.
25. De Clerq K, Mertens P, De Leeuw I, Oura C, Houdart P, Potgieter AC, Maan S, Hooyberghs J, Batten C, Vandemeulenbroucke E *et al.*: **Emergence of bluetongue serotypes in Europe, part 2: the occurrence of a BTV-11 strain in Belgium**. *Transbound Emerg Dis* 2013, **56**:355-361.
26. Reviriego Gordejo FJ: **Bluetongue in Europe: lessons learned**. *Proc US Anim Health Assoc* 2013, **117**:141-142.
27. Orłowska A, Trebas P, Smreczak M, Marzec A, Żmudziński JF: **First detection of bluetongue virus serotype 14 in Poland**. *Arch Virol* 2016, **161**:1969-1972.

This study describes a genetically novel strain of bluetongue virus that was introduced in sheep from Libya to Tunisia. The virus is genetically related to one recently identified as a contaminant of a commercial sheep pox vaccine. This finding further highlights the potential for bluetongue virus, including genetically novel strains, to contaminate live attenuated vaccines.

28. McVey DS, MacLachlan NJ: **Vaccines for prevention of bluetongue and epizootic hemorrhagic disease in livestock: a North American perspective**. *Vector Borne Zoonotic Dis* 2015, **15**:385-396.

A recent comprehensive review of the advantages and inherent deficiencies of existing orbivirus vaccines, and analysis of future needs and directions for improved vaccines.

29. Hofmann MA, Renzullo S, Mader M, Chaignat V, Worwa G, Thuer B: **Genetic characterization of Toggenburg Orbivirus, a new bluetongue virus, from goats, Switzerland**. *Emerg Infect Dis* 2008, **14**:1855-1861.
30. Chaignat V, Worwa G, Scherrer N, Hible M, Ehrensperger F, Batten C, Cortyen M, Hofmann M, Thuer B: **Toggenburg**

- Orbivirus, a new bluetongue virus: initial detection, first observations in field and experimental infection of goats and sheep.** *Vet Microbiol* 2009, **138**:11-19.
31. Casaubon J, Chaignat V, Vogt HR, Michel AO, Thuer B, Ryser-Deggorgis MP: **Survey of bluetongue virus infection in free ranging wild ruminants in Switzerland.** *BMC Vet Res* 2013, **9**:166.
 32. Planzer J, Kaufmann C, Worwa G, Gavier-Widen D, Hofmann MA, Chaignat V, Thuer B: **In vivo and in vitro propagation and transmission of Toggenburg orbivirus.** *Res Vet Sci* 2011, **91**: e163-e168.
 33. Marcacci M, Sant S, Mangone I, Gorla M, Dondo A, Zoppi S, van Gennip RGP, Radaelli MC, Cammà C, van Rijn PA *et al.*: **One after the other: a novel Bluetongue virus strain related to Toggenburg virus detected in the Piedmont region (North-western Italy), extends the panel of novel atypical BTV strains.** *Transbound Emerg Dis* 2018, **65**:370-374.
- The study describes a novel strain of bluetongue virus in the Piedmont region of Italy. Ruminants infected with this virus displayed low levels of RNAemia and absence of clinical signs, a scenario described in animals infected with other atypical strains of bluetongue virus.
34. Maan S, Maan NS, Nomikou K, Batten C, Antony F, Belaganahali MN, Samy AM, Reda AA, Al-Rashid SA, El Batel M *et al.*: **Novel bluetongue virus serotype from Kuwait.** *Emerg Infect Dis* 2011, **17**:886-889.
 35. Batten C, Henstock MR, Bin-Tarif A, Steedman HM, Waddington S, Edwards L, Oura CA: **Bluetongue virus serotype 26: infection kinetics and pathogenesis in Dorset poll sheep.** *Vet Microbiol* 2012, **157**:119-124.
 36. Batten C, Henstock MR, Steedman HM, Waddington S, Edwards L, Oura CA: **Bluetongue virus serotype 26: infection kinetics, pathogenesis and possible contact transmission in goats.** *Vet Microbiol* 2013, **162**:62-67.
 37. Lorusso A, Bada D, Spedicato M, Teodori L, Bonfini B, Marcacci M, Di Provido A, Isselmou K, Marini V, Carmine I *et al.*: **Bluetongue surveillance in the Islamic Republic of Mauritania: is serotype 26 circulating among cattle and dromedaries?** *Infect Genet Evol* 2016, **40**:109-112.
 38. Zientara S, Sailleau C, Vigarouge C, Hoper D, Beer M, Jenekel M, Hoffmann B, Romey A, Bakkali-Kassimi L, Fablet A *et al.*: **Identification of a novel bluetongue virus in goats, Corsica, France, 2014.** *Transbound Emerg Dis* 2014, **20**:2123-2125.
 39. Bréard E, Schulz C, Sailleau C, Bernelin-Cottet C, Viarouge C, Vitour D, Guillaume B, Caignard G, Gorlier A, Attoui H *et al.*: **Bluetongue virus serotype 27: experimental infection of goats, sheep and cattle with three BTV-27 variants reveal atypical characteristics and likely direct contact transmission BTV-27 between goats.** *Transbound Emerg Dis* 2018, **65**:e251-e263.
 40. Savini G, Puggioni G, Meloni G, Marcacci M, Di Domenico M, Rocchigiani AM, Spedicato M, Oggiano A, Manunta D, Teodori L *et al.*: **Novel putative Bluetongue virus in healthy goats from Sardinia, Italy.** *Infect Genet Evol* 2017, **51**:108-117.
 41. Sun EC, Huang LP, Xu QY, Wang HX, Xue XM, Lu P, Li WJ, Liu W, Bu ZG, Wu DL: **Emergence of a novel bluetongue virus serotype, China 2014.** *Transbound Emerg Dis* 2016, **63**:585-589.
- This study confirms the presence of novel bluetongue viruses beyond Europe and the Middle East, emphasizing the need for ongoing surveillance.
42. Bumbarov V, Golender N, Erster O, Khinich Y: **Detection and isolation of Bluetongue virus from commercial vaccine batches.** *Vaccine* 2016, **34**:3317-3323.
 43. Belbis G, Zientara S, Breard E, Sailleau C, Caignard G, Vitour D, Attoui H: **Bluetongue virus: from BTV-1 to BTV-27.** *Adv Virus Res* 2017, **99**:161-197.
 44. Akita GY, Ianconescu M, MacLachlan NJ, Osburn BI: **Bluetongue disease in dogs associated with contaminated vaccine.** *Vet Rec* 1994, **134**:283-284.
 45. Wilbur LA, Evermann JF, Levings RL, Stoll IR, Starling DE, Spillers CA, Gustafson GA, McKeirnan AJ: **Abortion and death in pregnant bitches associated with a canine vaccine contaminated with bluetongue virus.** *J Am Vet Med Assoc* 1994, **204**:1762-1765.
 46. Dubovi EJ, Hawkins M, Griffin RA Jr, Johnson DJ, Ostlund EN: **Isolation of Bluetongue virus from canine abortions.** *J Vet Diagn Invest* 2013, **25**:490-492.
 47. Gaudreault NN, Jaspersen DC, Dubovi EJ, Johnson DJ, Ostlund EN, Wilson WC: **Whole genome sequence analysis of circulating Bluetongue virus serotype 11 strains from the United States including two domestic canine isolates.** *J Vet Diagn Invest* 2015, **27**:442-448.
 48. Oura CA, El Harrak M: **Midge-transmitted bluetongue in domestic dogs.** *Epidemiol Infect* 2011, **139**:1396-1400.
 49. O'Dell N, Arnot L, Janisch CE, Steyl JC: **Clinical presentation and pathology of suspected vector transmitted African horse sickness in South African domestic dog from 2006 to 2017.** *Vet Rec* 2018, **182**:715.
 50. Osborne CJ, Mayo CE, Mullens BA, McDermott EG, Gerry AC, Reisen WK, MacLachlan NJ: **Lack of evidence for laboratory and natural vertical transmission of Bluetongue virus in *Culicoides sonorensis* (Diptera: Ceratopogonidae).** *J Med Entomol* 2015, **52**:274-277.
 51. Mayo CE, Mullens BA, Reisen WK, Osborne CJ, Gibbs EP, Gardner IA, MacLachlan NJ: **Seasonal and interseasonal dynamics of bluetongue virus infection of dairy cattle and *Culicoides sonorensis* midges in northern California—implications for virus overwintering in temperate zones.** *PLoS One* 2014, **9**:e106975.
 52. Mayo C, Mullens B, Gibbs EP, MacLachlan NJ: **Overwintering of Bluetongue virus in temperate zones.** *Vet Ital* 2016, **52**:243-246.
 53. McDermott EG, Mayo CE, Gerry AC, Mullens BA: **Trap placement and attractant choice affect capture and create sex and parity biases in collection of the biting midge, *Culicoides sonorensis*.** *Med Vet Entomol* 2016, **30**:293-300.
 54. McDermott EG, Mayo CE, Gerry AC, Laudier D, MacLachlan NJ, Mullens BA: **Bluetongue virus infection creates light averse *Culicoides* vectors and serious errors in transmission risk estimates.** *Parasites Vectors* 2015, **8**:460.
- This study confirms for the first time that exclusive use of light-trapping to estimate transmission risk of bluetongue virus is likely to lead to serious errors in risk assessment. Specifically, the study confirms generalized bluetongue virus infection in adult female midges results in disseminated infection, including of nervous tissues and eyes, which may result in unanticipated behavioral effects including attraction to light traps.
55. Golender N, Khinich Y, Gorohov A, Abramovitz I, Bumbarov V: **Epizootic hemorrhagic disease virus serotype 6 outbreak in Israeli cattle in 2015.** *J Vet Diagn Invest* 2017, **29**:885-888.
 56. Kamomae Y, Kamomae M, Ohta Y, Nabe M, Kagawa Y, Ogura Y, Kato T, Tanaka S, Yanase T, Shirafuji H: **Epizootic hemorrhagic disease virus serotype 6 infection in cattle, Japan, 2015.** *Emerg Infect Dis* 2018, **24**:902-905.
 57. Hassine TB, Amdouni J, Monaco F, Savini G, Sghaier S, Selimen IB, Chandoul W, Hamida KB, Hammami S: **Emerging vector-borne diseases in dromedaries in Tunisia: West Nile, bluetongue, epizootic haemorrhagic disease and Rift Valley fever.** *Onderstepoort J Vet Res* 2017, **84**:e1-e3.
 58. MacLachlan NJ, Drew CP, Darpe KE, Worwa G: **The pathology and pathogenesis of bluetongue.** *J Comp Pathol* 2009, **141**:1-16.
 59. MacLachlan NJ, Osburn BI: **Teratogenic bluetongue and related orbivirus infections in pregnant ruminant livestock: timing and pathogen genetics are critical.** *Curr Opin Virol* 2017, **27**:31-35.
 60. Feenstra F, van Rijn PA: **Current and next-generation bluetongue vaccines: requirements, strategies, and prospects for different field situations.** *Crit Rev Microbiol* 2017, **43**:142-155.
 61. Mayo C, Lee J, Kopanke J, MacLachlan NJ: **A review of potential bluetongue virus vaccine strategies.** *Vet Microbiol* 2017, **206**:84-89.
 62. Noad R, Roy P: **Bluetongue vaccines.** *Vaccine* 2009, **27**(Suppl. 4):D86-D89.